## **Compounds in Aqueous Solutions**

## When ionic compounds dissolve in water, it is called *dissociation* (already in your notes)

 $NaCl(s) + H_2O(l)$   $Na^+(aq) + Cl^-(aq)$ 

Are all ionic compounds soluble in water??

7 General solubility guidelines

1. All salts of Group IA, and ammonium are soluble.

2. All salts of nitrates, chlorates and acetates are soluble.

3. All salts of halides are soluble except those of silver(I), copper(I), lead(II), and mercury(I).

4. All salts of sulfate are soluble except for barium sulfate, lead(II) sulfate, and strontium sulfate.

5. All salts of carbonate, phosphate and sulfite are insoluble, except for those of group IA and ammonium.

6. All oxides and hydroxides are insoluble except for those of group IA, calcium, strontium and barium.

7. All salts of sulfides and insoluble except for those of Group IA and IIA elements and of ammonium.

**Determining precipitates** 

When a double replacement occurs, determine if one of the products is insoluble.

-write a balanced equation for ammonium sulfide and cadmium nitrate and determine the phase for each product (both reactants are aq)

Net Ionic Equations

an equation in which only those compounds and ions that undergo a chemical change in a reaction in an aqueous solution

Start with the ionic version of the balanced equation...

 $Cd^{2+}(aq) + 2NO_{3}(aq) + 2NH_{4}(aq) + S^{2}(aq) \longrightarrow$ 

 $CdS(s) + 2NO_{3}(aq) + 2NH_{4}(aq)$ 

Notice that some ions are the same on both sides of the equation. These ions did not undergo any chemical change, hence they are called *spectator ions*. To write the **net ionic equation**, simply remove the ones that are the same on both sides.

 $Cd^{2+}(aq) + S^{2-}(aq) \longrightarrow CdS(s)$ 

Ionization

(already in your notes) is when a covalent molecule separates into ions in a solution

HCl water, H+(aq) + Cl-(aq)

Many molecular compounds have a H bonded by a polar covalent bond, when these H ionize they then react with the water to form the *hydronium ion* 

 $H_2O(I) + HCI(g) \longrightarrow H_3O^+(aq) + CI^-(aq)$ 

The formation of the hydronium ion releases energy which helps to break apart more the molecular solute

## Strong Electrolytes

-any compound whose dilute aqueous solutions conduct electricity well; this is due to the presence of all or almost all of the dissolved compound in the form of ions.

examples: HCI, HBr, HI, several other acids, and all soluble ionic compounds


Weak Electrolytes

-any compound whose dilute aqueous solutions conduct electricity poorly; this is due to the presence of a small amount of the dissolved compound in the form of ions

 $HF(aq) + H_2O \longrightarrow H_3O^+(aq) + F^-(aq)$ 

**Colligative Properties of Solutions** 

-properties that depend on the concentration of solute particles but not on their identity



**Freezing-Point Depression** 

molal freezing point constant for water is

 $K_{f} = -1.86 \text{ K/}m$ 

K<sub>f</sub> is different for different solvents.

freezing point depression is how much the freezing point is lowered due to the concentration of particles of solute in a solvent -

 $\Delta t_f = K_f m$ 

**Boiling-Point Elevation** 

molal boiling point constant for water is

 $K_{b} = 0.51 \text{ K/m}$ 

K<sub>b</sub> is different for different solvents.


boiling point elevation is how much the boiling point is raised due to the concentration of particles of solute in a solvent -

 $\Delta t_{b} = K_{b}m$ 

Osmotic Pressure

**osmosis** is the movement of solvent through a semipermeable membrane from the side of lower solute concentration tot he side of higher solute concentration

**osmotic pressure** is the external pressure that must be applied to stop osmosis



**Electrolytes and Colligative Properties** 

-when nonelectrolytes dissolve, they particles remain whole, so a 1 *m* solution of sucrose has 1 mol of particles in it. When electrolytes dissolve, the number of particles increases.

-NaCl disassociates into Na and Cl ions, hence 1 *m* NaCl becomes 2 mols of particles. 1 *m* CaCl<sub>2</sub> becomes 3 mols of particles.

Determine the freezing point depression of 62.5 g Ba(NO<sub>3</sub>)<sub>2</sub> dissolved in 1.00 kg of water.  $K_f H_2O = -1.86K$ 

7-Quiz 12 Molarity-Molality, Solutions and Colligative Properites.notebook February 24, 2020

Hydrogen sulfide is bubbled through a solution of strontium hydroxide

A manganese (II) nitrate solution is mixed with a sodium hydroxide solution

Solutions of sodium carbonate and lead (II) nitrate are mixed